

Journal of Integrative Medicine

Volume 20, Issue 4, July 2022, Pages 284–287

Commentary

The scope and potential of music therapy in stroke rehabilitation

Tara Rajendran ^a ✉ ... Martha Summa-Chadwick ^b

Show more

Share Cite

<https://doi.org/10.1016/j.joim.2022.04.006>

Get rights and content

Abstract

There is a growing interest in the use of music therapy in neurological rehabilitation. Of all the major neurological illnesses, stroke rehabilitation has been observed to have some of the strongest potential for music therapy's beneficial effect. The current burden of stroke has raised the need to embrace novel, cost-effective, rehabilitation designs that will enhance the existing physical, occupation, and speech therapies. Music therapy addresses a broad spectrum of motor, speech, and cognitive deficits, as well as behavioral and emotional issues. Several music therapy designs have focused on gait, cognitive, and speech rehabilitation, but most of the existing randomized controlled trials based on these interventions have a high risk of bias and are statistically insignificant. More randomized controlled trials with greater number of participants are required to strengthen the current data. Fostering an open and informed dialogue between patients, healthcare providers, and music therapists may help increase quality of life, dispel fallacies, and guide patients to specific musical interventions.

Introduction

Stroke is the third leading universal cause of death and disability [1]. A third of stroke survivors are dependent on others for their care; increasing number of stroke survivors create greater demand for novel interventions that interact with the mechanisms of spontaneous biological recovery [2], [3]. Rehabilitation is one of the critical components of patient care that mitigate post-stroke disability and dependency [4]. Stroke rehabilitation comprises a wide range of strategies, including skilled therapy interventions that address mobility and activities of daily living, evaluation and treatment of communication and cognitive impairments, and treatment of dysphagia [3], [4]. The current burden of stroke has raised the need to embrace novel, cost-effective, rehabilitation designs that will enhance the existing physical, occupation, and speech therapies [5]. Until the late 1990s, stroke rehabilitation was criticized for a lack of scientific evidence base [2]. However, the last two decades of scientific investigations on potential adjunct rehabilitation designs have positively changed this discernment [5]. Of all the major neurological illnesses, stroke rehabilitation has been observed to have some of the strongest potential for music therapy's beneficial effect [5]. Emerging randomized controlled trial (RCT) findings suggest that music therapy holds promise for stroke-related neurologic and neuropsychiatric impairments, including motor rehabilitation, speech regeneration, and cognitive recovery [6], [7], [8], [9], [10], [11], [12], [13], [14], [15], [16], [17], [18], [19], [20], [21].

Section snippets

Brain plasticity, music, and stroke rehabilitation

Brain plasticity refers to brain's innate ability to reorganize its function and structure in response to stimuli and injuries [22]. Following stroke, the plasticity process commences in an attempt to compensate for the lesion as well as for the remote effects. These changes may take place in the days, months, and years following the stroke [22], [23]. Researchers suggest utilizing this time-limited window of brain plasticity as this leads to the greatest gain in recovery [24]. Various...

Music therapy and neurologic music therapy

Music therapy is the clinical and evidence-based use of music interventions to accomplish individualized goals within a therapeutic relationship, by a credentialed professional who has completed an approved music therapy program [38], [39]. Neurologic music therapy is a domain of music therapy where evidence-based and standardized therapeutic music interventions address sensorimotor, cognitive, and speech and language dysfunctions resulting from neurological illnesses and impairments, as well...

Challenges and recommendations

Guidelines for Adult Stroke Rehabilitation and Recovery are developed by the American Heart Association/American Stroke Association [55]. In order to be recommended by the guidelines, music-based interventions require strong evidence in the form of large-scale rigorous clinical trials in multiple populations [55]. While RCTs report encouraging results for the effects of various music therapy interventions on several parameters, the quality of many RCTs on musical intervention in stroke...

Conclusion

Stroke rehabilitation is often multidimensional and requires an interdisciplinary, comprehensive, patient-centric and patient-directed approach to management and rehabilitation. Exploring the potential benefits of music is essential for advancing clinical practice. A genuine partnership between music therapists, speech therapists and physicians can draw on the strengths of each to benefit research and improve clinical practice. Patient experience will be far more positive if clinical practice...

Funding

No funding was received for this study...

Authors' contribution

TR and MSC did the conceptualization, drafting, and editing of the article...

Declaration of competing interests

The authors declare that they have no competing interests...

References (60)

T. Rajendran
Addressing the need for personalizing music therapy in integrative oncology
J Integr Med (2022)

S.V. Norman-Haignere *et al.*
A neural population selective for song in human auditory cortex
Curr Biol (2022)

M.H. Thaut
The discovery of human auditory-motor entrainment and its role in the development of neurologic music therapy
Prog Brain Res (2015)

V. Menon *et al.*
The rewards of music listening: response and physiological connectivity of the mesolimbic system
Neuroimage (2005)

M.L. Chanda *et al.*
The neurochemistry of music
Trends Cogn Sci (2013)

E. Altenmüller *et al.*
Apollo's gift: new aspects of neurologic music therapy
Prog Brain Res (2015)

M.H. Thaut *et al.*
Rhythmic facilitation of gait training in hemiparetic stroke rehabilitation
J Neurol Sci (1997)

S. Jeong *et al.*
Effects of a theory-driven music and movement program for stroke survivors in a community setting
Appl Nurs Res (2007)

A.J. Sihvonen *et al.*
Music-based interventions in neurological rehabilitation
Lancet Neurol (2017)

C.M. Stinear *et al.*
Advances and challenges in stroke rehabilitation
Lancet Neurol (2020)

View more references

Cited by (0)

Recommended articles (6)

Research article
Remission of type 2 diabetes depends on prompt comprehensive lifestyle changes upon diagnosis: How can this “Road to Damascus” experience be supported?
Journal of Integrative Medicine, Volume 20, Issue 4, 2022, pp. 288–291
Show abstract

Research article
Can acupuncture enhance therapeutic effectiveness of antidepressants and reduce adverse drug reactions in patients with depression? A systematic review and meta-analysis
Journal of Integrative Medicine, Volume 20, Issue 4, 2022, pp. 305–320
Show abstract

Research article
Ziyin Huatan Recipe, a Chinese herbal compound, inhibits migration and invasion of gastric cancer by upregulating RUNX3 expression
Journal of Integrative Medicine, Volume 20, Issue 4, 2022, pp. 355–364
Show abstract

Research article
To “tell or not to tell”—Exploring disclosure about medicine use by people living with sleep disorders
Journal of Integrative Medicine, Volume 20, Issue 4, 2022, pp. 338–347
Show abstract

Research article
Prevalence and Predictors of Successful Percutaneous Coronary Intervention in Ostial Chronic Total Occlusion
JACC: Cardiovascular Interventions, Volume 15, Issue 10, 2022, pp. 1094–1096

Research article
Prevalence of long term metabolic, cardiovascular, cerebrovascular and renal disease in patients with hypertensive disorders in pregnancy remote from pregnancy (POMCH)
Pregnancy Hypertension, Volume 28, 2022, pp. 162–167
Show abstract

View full text